



Nutritional Rehabilitation Care of Refeeding Syndrome with Anorexia Nervosa

Tara Khursheed^{1*}, Shahnai Bashrat², Madiha Jaffar² and Hafsa Tahir²

¹Department of Diet and Nutritional Sciences, The University of Lahore, Pakistan

²Department of Diet and Nutritional Sciences, University of Management and Technology, Pakistan

Abstract

The term “Refeeding syndrome” is defined by the immense changes in the electrolytes and fluid levels that could take place in the patients who are refeed artificially after suffering a long term malnourishment. An improved level of awareness as well as understanding of refeeding syndrome, with effectively managed strategy is important for nutritional rehabilitation of the individual. In this scenario, a gradual and deliberate surge in nutritional intake with proper nutritional counseling, and psychotherapy are recommended by the experts. Our case study took a young 22 year-old girl suffering from anorexia with a BMI of 11.9. She had been misdiagnosed by the medical practitioner and got a severe electrolyte imbalance in blood levels. This occurred only because of the lack of cognizance regarding refeeding syndrome. After the cautious clinical and nutritional assessment, empirical and scientific evaluation, different protocols can be suggested to prevent and manage this syndrome. Equally important is to instill awareness in the patient about refeeding slowly and building-up micronutrient gradually for several days in nutritional therapy.

Keywords: Refeeding syndrome; Malnourishment; Anorexia nervosa; Nutritional rehabilitation

Introduction

Refeeding Syndrome (RS) is possibly a serious condition, that is defined by severe fluid and electrolyte imbalances cause by damaging nutritional rehabilitation in patients who are conditioned to malnourishment [1]. The patients who are enduring protein energy malnutrition and severe weight loss are at a potential risk of developing RS in them. Those having condition such as anorexia nervosa and obesity are at high end risk [2]. Even though there are studies but the actual incidence of refeeding syndrome is still unknown to us. However a study on the topic of RS has demonstrated a prevalence rate of 0.45% in all of the hospitalized patients, whereas malnutrition is one of the major risk factor [3]. The principal features of refeeding syndrome comprise severe hypophosphatemia, hypo-magnesia, and hypokalemia with other fluid imbalances and electrolyte abnormalities that could develop life-threatening irregularities in the central nervous system, cardiovascular system, including different pulmonary and hematologic complications [4]. Refeeding malnourished patients can emerge from “oral, enteral and parenteral nutrition” which in several cases could be life threatening for the patient if it goes undiagnosed and ill-treated.

The incident of a 22-year old girl who had previously been suffering from the condition of anorexia nervosa is well known. She developed refeeding syndrome due an aggressive nutrition intake and was managed in a hospital with “Total Parenteral Nutrition” (TPN).

Case Presentation

The 22-year-old girl living in Lahore had been admitted in the emergency care unit on September, 2021. She had a 2 days history of severe abdominal cramps, nausea and continuous vomiting. She endured the condition of secondary amenorrhea, and chronic constipation for two and half years before admission to the hospital. Notwithstanding, she had a psychiatric history of having anorexia nervosa, which is the fear of growing overweight and was on strict dieting schedule. The patient was asked to consume a large serving of meat just a day before her admission to hospital. She instantly developed afore-mentioned symptoms after eating the meal. Over the course fourteen weeks she lost eighteen kg of her weight. Upon examination, her weight was only twenty seven kilo grams and was 152 cm tall. As her Body Mass Index (BMI) stood at 11.9 which is extremely low. In addition to this her physical appearance was pale and weak. Her resting pulse rate was rapid (110 beats/minute), with a low blood pressure measuring just 90/40 mm/Hg. Upon her Chest and CVS examination

OPEN ACCESS

*Correspondence:

Tara khursheed, Department of Diet and Nutritional Sciences, The University of Lahore, Pakistan,
E-mail: tarakhursheed93@gmail.com

Received Date: 28 Jan 2022

Accepted Date: 15 Feb 2022

Published Date: 21 Feb 2022

Citation:

Khursheed T, Bashrat S, Jaffar M, Tahir H. Nutritional Rehabilitation Care of Refeeding Syndrome with Anorexia Nervosa. *Clin Case Rep Int.* 2022; 6: 1289.

Copyright © 2022 Tara Khursheed.

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1: Biochemical values.

Biochemical values	Admission values	Reference values
Hb (g/dl)	12.4	13.5-16.3
Platelets (× 10 ⁹ /l)	145	150-400
Albumin (g/dl)	2.1	3.5-4.8
Phosphate (mmol/l)	1.5	2.2 -4.8
Potassium (mmol/l)	2.4	3.6-5
Magnesium (mmol/l)	1.6	1.9-2.5
Calcium (mmol/l)	6.6	8-11
Chloride (mmol/l)	72	104-114
Sodium (mmol/l)	122	136-138
BUN (mg/dl)	11	6-16
Creatinine (mg/dl)	0.7	0.85-1.35
Bicarbonate (mEq/l)	34.6	17.5-27.5
Total bilirubin (mg/dl)	5.5	0.25-1
SGPT (IU/L)	146	03-33
SGOT (IU/L)	172	18-32

nothing abnormal was found. However, the abdominal examination of the patient showed recessed abdomen with a moderate tenderness in her right iliac fossa and the visceral organs were not palpable. The laboratory tests of the girl revealed low albumin, hypophosphatemia, hypochloremic metabolic acidosis, hypokalemia. Her liver function tests showed a disturbed liver function (Table 1). She had turn to several gynecologists during the last 2 years to consult secondary amenorrhea. She was recommended some medicines to regulate her

Table 2: Clinical signs and symptoms of refeeding syndrome [1].

Hypophosphatemia	Hypokalemia	Hypomagnesia	Thiamine deficiency	Na retention	Hyperglycemia
Dyspnea	Arrhythmias	Arrhythmias	Encephalopathy	Edema	Hypotension
Hypotension,	Muscle necrosis	Tremors	Lactic acidosis	Fluid overload	Respiratory failure
Respiratory failure	Severe Constipation	Seizures	-	-	Metabolic ketoacidosis
Shock	Rhabdomyolysis	Coma	-	-	Dehydration
Arrhythmias	-	Diarrhea	-	-	-
Cardiac failure	-	-	-	-	-

Table 3: Guidelines for replacement therapy [1].

Phosphate	Intravenous replacement dose (administer over 6-12 hours)
	Children: 0.07-0.23 mmol/kg
	Maximum daily dose: 1.5 mmol/kg
	Adults: Initially 0.08-0.09 mmol/kg for mild and 0.17 mmol/kg for severe hypophosphatemia
	Maximum dose: 0.23-0.24 mmol/kg per dose
Magnesium	Intravenous replacement dose (administer over 4-5 hours)
	Children: 25-50mg/kg per dose (0.2-0.5 mEq/kg per dose)
	Maximum single dose: 2000 mg (16 mEq)
	Adults: 1 g every 6 hours for 4 doses for moderate and 8-12 g/d in divided doses for severe
Potassium	Intravenous replacement dose (administer over 1 hr)
	Children: 0.3 - 0.5 mEq/kg per dose
	Max Dose: 30 mEq per dose
	Adults: same
Thiamine	Children: 10-25 mg/d administered PO everyday for 2 weeks and then 5-10 mg/d for 1 month.
	Adults: 5-31 mg per dose 3 times per day, administered IV and then 5-31 mg/d administered PO for one month

monthly cycles, but her prolactin levels elevated later on. She was pre-occupied with the feelings of being overweight, despite the fact that her weight was only 42 kg with BMI of 18.6. The fear lead her to stop taking her medicines and she started dieting and vigor exercise. Afterwards she was taken for a psychiatrist consultation and there she was diagnosed with the condition of anorexia nervosa and was advised to take anti-depressants. In her case, no family psychiatric history was found and there was no major childhood illness. She was again presented to hospital 3 days later with stark phosphatemia, abdominal distension, feet swelling. There she was diagnosed with refeeding syndrome as her reports were claiming severe electrolyte imbalances. Then she was recommended for nutritional consultation was given TPN 1 (for precise intake). Later on her schedule was changed to TPN 2 as she was unable tolerate the oral intake. She had been prescribed a 1,500 kcal/day diet plan and antiemetics before meals to stop nausea and vomit. She was also tied up into psychotherapy and motivational therapy. There was steady improvement in her condition and she was gradually substituted to a total oral consumption. In her 5th week of therapy, she added 6 kg in her total weight. She was discharged from hospital with a prescription of esomeprazole, some antidepressants and other vitamin supplements.

Discussion

A forceful nutrient intake of carbs can evoke a release of insulin and increase levels of glucose, phosphate, magnesium etc in the blood. The severe electrolyte imbalances can end in severe complications including sudden death of the patients. Here, the natriuretic effect of insulin can cause sodium retention and edema. Severe phosphatemia is the most prevailing feature of refeeding syndrome and can cause “dyspnea, respiratory failure, hypotension, shock, arrhythmias and

cardiac failure". The increase in the utilization of thiamine in such conditions can be resulted into thiamine deficiency in some of the patients, as indicated by the study of Wernicke's encephalopathy [5]. Other clinical symptoms are described in Table 2.

A procedure for avoidance and controlling of refeeding syndrome specified the circumstances in which it is probably develop, and stressed the significance of refeeding gradually and compounding the micro nutritional content over few days with constant monitoring [4]. It highlighted that the administration should never be performed in haste and the targeted weight accumulation should never surpass 1 kg to 1.5 kg per week. In the adults, studies suggested the starting the intake of 1000 to 1200 kcal per day [6]. The protein intake must not be circumscribed during the nutritional provision [7]. The strategies for electrolyte and thiamine stabilization are given in the Table 3.

In conclusion, the 'nutritional rehabilitation' is always a crucial element of the physical treatment of the disorders such as 'anorexia nervosa' and other conditions that may cause malnourishment. However, good psychological treatment is also of paramount importance in order to address the anxiety issues concerning the patient. An upright therapeutic coalition between the doctor, patient and the family creates a collaboration that allows the work of motivational factors. The treatment at home is conferred in some studies is found to bear more benefits for the patients [1].

So, improved awareness and understanding of refeeding syndrome along with a well-coordinated plan are important in delivering safe and effective nutritional rehabilitation.

In summary, rehabilitation of anorexia nervosa or other starving states can result in fatal electrolyte disturbances (refeeding syndrome) and, therefore, proper and timely nutrition is recommended. We suggest a slow and gradual increase in nutrition along with nutritional counseling and psychotherapy, including the patient's family, to prevent resistance from the patient and to ensure long-term compliance by the patient.

References

1. Fuentebella J, Kerner JA. Refeeding syndrome. *Pediatr Clin North Am.* 2009;56(5):1201-10.
2. Khan LU, Ahmed J, Khan S, MacFie J. Refeeding syndrome: A literature review. *Gastroenterol Res Pract.* 2011;2011:410971.
3. Mehanna H, Nankivell PC, Moledina J, Travis J. Refeeding syndrome—awareness, prevention and management. *Head Neck Oncol.* 2009;1(1):4.
4. Stanga Z, Brunner A, Leuenberger M, Grimble RF, Shenkin A, Allison SP, et al. Nutrition in clinical practice—the refeeding syndrome: Illustrative cases and guidelines for prevention and treatment. *Eur J Clin Nutr.* 2008;62(6):687-94.
5. Reuler JB, Girard DE, Cooney TG. Wernicke's encephalopathy. *NEJM.* 1985;312(16):1035-9.
6. Miller SJ. Death resulting from overzealous total parenteral nutrition: The refeeding syndrome revisited. *Nutr Clin Pract.* 2008;23(2):166-71.
7. Palesty JA, Dudrick SJ. The goldilocks paradigm of starvation and refeeding. *Nutr Clin Pract.* 2006;21(2):147-54.